

The prevalence and predisposing factors of mastitis in lactating mothers in puerperium

Ruma Dutta, Ravikanth Onkarappa Gowder

Correspondence: Ravikanth Onkarappa Gowder, Associate Professor, Dept. of Obstetrics and Gynaecology, K.V.G Medical College and Hospital, Sullia, Rajiv Gandhi University Of Health Sciences, Karnataka, India; Email - drravikanthgo@gmail.com

Distributed under Creative Commons Attribution-Share Alike 4.0 International.

ABSTRACT

Objectives: To study the prevalence, evaluate the predisposing factors and know the etiological factors associated with mastitis in lactating mothers during puerperium. **Materials and methods:** A prospective study was performed from January 1st, 2014 to December 31st, 2015. Data were collected from patients admitted in the postnatal ward with benign breast problems by preformatted questionnaires. Breast milk cultures were done to evaluate the etiological factors. Data were analyzed by using descriptive statistics. **Results:** The overall prevalence of puerperal mastitis among lactating mothers was 5.1%, high among caesarean section patients (5.8%) compared to vaginally delivered patients (4.8%). The predisposing factors include primipara (57%), unbooked cases (57%), low socioeconomic status (49%), anemic patients (45%), breast engorgement (75%), cracked nipple (44%), retracted/flat/inverted nipple (13%), poor attachment of baby to breast (32%), infrequent removal of milk (7%), oversupply of breast milk (37.5%) and lower supply of breast milk (12.5%). Breast milk culture reports yielded growth of *Staphylococcus aureus* (75%) and MRSA (25%). Seven patients developed breast abscess (4.4%). **Conclusion:** It is recommended to educate the mothers and the family on the exclusive breast feeding practice and its benefits, demand feeding, optimal positioning and attachment of the baby, to provide adequate emotional support and access to skilled help in the early postpartum period.

Keywords: Puerperium, mastitis, lactation.

Mastitis is defined as inflammation of the mammary gland. Mastitis is an acute, debilitating condition that occurs in approximately 20 % of breastfeeding women who experience a red, painful breast with fever^{1, 2}. Two mode of infection, firstly involving the parenchymatous breasts tissues which may lead to cellulitis. The lacteal system remains unaffected. Secondly, infection gains access through the lactiferous duct leading to development of primary mammary adenitis. Almost always from nursing infant's nose and throat, the

organism enters the breast through the nipple at the site of a fissure or abrasion. In superficial cellulitis, the onset is acute during first 2-4 weeks postpartum. There may be inflammation preceded by engorgement, fever (102°F or more) with chills, tachycardia, flu like symptoms like generalized malaise, headache, nausea, vomiting, sudden onset of intense breast pain. Breast abscess usually occurs as a complication of mastitis³.

Materials and Methods

This study was conducted among 3117 postnatal

Received: 24th January 2018. **Accepted:** 17th April 2018.

Dutta R, Gowder RO. The prevalence and predisposing factors of mastitis in lactating mothers in puerperium. The New Indian Journal of OBGYN. 2018; 5(1): 28-32.

lactating mothers who had a full term or preterm, vaginal or caesarean delivery at K.V.G medical college & Hospital, Sullia, Karnataka, from January 1st 2014 to December 31st 2015, for a period of six weeks postpartum. Out of 3117 number of delivery, 160 mothers developed mastitis. Data were collected from patients admitted in the postnatal ward with benign breast problems like cracked nipple or sore nipple, breast engorgement, retracted nipple, flat nipple or inverted nipple, poor attachment of baby to the breasts leading to nipple damage, infrequent removal of milk, breast milk amount (normal/low supply/over supply) and breast abscess by preformatted questionnaires and local examination of breasts. Breast milk cultures were done to evaluate the etiological factors. The exclusion criteria of our study were history of mastitis prior to delivery and continued in the postnatal period. Data were analyzed by using descriptive statistics and other appropriate statistical methods.

Results

The overall prevalence of puerperal mastitis in lactating mothers in this study was 5.1%. Out of 2084 vaginally delivered patients, 100 patients developed puerperal lactational mastitis, the incidence was 4.8%.

Table 1: Demographic Data (N=160)

Categories	Number (%)
Booked patients	69(43%)
Unbooked patients	91(57%)
Primipara	91(57%)
Age in years	
18-23	45(28%)
24-30	88(55%)
>30	27(17%)
Mean age	27 yrs; range 18-36yrs
Socioeconomic status (According to Kuppuswamy classification)	
Upper	13(8%)
Upper middle	5(3%)
Lower middle	64(40%)
Upper lower	78(49%)

Out of 1033 patients, who underwent cesarean section, 60 patients developed puerperal lactational mastitis, the incidence was 5.8%. Among the 160 puerperal lactational mastitis patients, primipara were 91(57%) and multipara were 69 (43%) (Table 1). The mean age of the women was 27 years (Range 18 to 36 years) (Table 1). The distribution of patients who developed puerperal mastitis

in relation to period of gestation <37weeks, 37- 40weeks and > 40weeks were 20(12.5%), 130 (81.25%) and 10(6.25%) respectively. In this study, lactational mastitis was found to be more in low socioeconomic status group (49%) and unbooked cases (57%) (Table 1).

The predisposing factors of lactational mastitis in this study was breast engorgement (75%), anemia (45%),

Table 2: Predisposing factors which might lead to puerperal mastitis among lactating mothers (N=160)

Risk factors	No of cases (%)
Primi parity	91(57%)
Low socio-economic status	78(49%)
Anemia	72(45%)
Breast engorgement	120(75%)
Cracked nipple	70(44%)
Retracted nipple	21(13%)
Poor attachment of baby to breast	51(32%)
Infrequent removal of milk	51(32%)
Breast milk amount	
Normal	80(50%)
Over supply	60(37.5%)
Low supply	20(12.5%)

cracked nipple (44%), poor attachment of baby to the breasts (32%), over supply of breast milk (37.5%) and lower supply of breast milk (12.5%) respectively (Table 2). Out of 72 anemic patients, 13 (8%) had mild, 40 (25%) had moderate and 19 (12%) patients had severe anemia.

Breast milk culture report of mastitis yielded growth of *Staphylococcus aureus* in 120 (75%) patients and MRSA (Methicillin resistance *Staphylococcus aureus*) in 40(25%) patients. The prevalence of breast abscess in our

Table 3: Breast milk culture report of patients

Breast milk culture	Mastitis (N=160)	Breast abscess (N=7)
	Number (%)	Number (%)
<i>Staphylococcus aureus</i>	120(75%)	5(71%)
MRSA	40(25%)	2(29%)

study was 4.4% (7/160) (Table 3). In one of the patients with breast abscess, one liter of pus which was drained with radial incision made over the breast. Breast Abscess milk culture yielded growth of *Staphylococcus aureus* and MRSA.

Discussion

Mastitis is defined as at least two breast signs or symptoms (pain, redness or lump) and one systemic

symptom (fever or 'flu-like symptoms) present for at least twelve hours⁴. Mastitis is reported to occur in 2-24% of breastfeeding women from several weeks up to one year after delivery in women who continue to breastfeed⁵. It is most common in the first 6 weeks of breastfeeding with the highest incidence occurring during the second and third weeks. It is initially localized to one segment of the breast, but if untreated, can spread to affect the whole breast^{6,7}.

We studied 3117 postnatal lactating mothers in puerperium, out of which 160 mothers developed mastitis. The prevalence of mastitis among lactating mothers in the first six weeks of postpartum period in the present study is 5.1%. The prevalence of mastitis varies depending on the definition and the number of weeks postpartum⁵. For example: A community based study among rural women of Rajasthan by Iyengar K reported the incidence of mastitis was 1.3% during first week after delivery⁵. A prospective cohort study conducted in western Nepal reported the incidence of mastitis was 8.0% in the first month postpartum⁸. The CASTLE study, a prospective cohort study which recruited nulliparous women in late pregnancy in two maternity hospitals in Melbourne, Australia reported 20% (70/346) of participants developed mastitis during the first eight weeks postpartum². The incidence rate of 9.5% (n = 946) was reported for a cohort of women followed up for 3 months in the United States⁸. A descriptive study of mastitis in Australian breastfeeding women reported a 6 month incidence rate of 17% (n = 206) of women experienced mastitis, of which 53% cases occurred in first 4 weeks postpartum¹. A prospective cohort study of mastitis among out of 670 Chinese breastfeeding mothers, 42 women (6.3%) experienced at least one episode of mastitis during the first 6 months after delivery^{8,9}. Scott et al. reported a 6-month incidence of 18% among a Scottish cohort, of which 53% of the cases (30 of 57) occurred in the first 4 weeks postpartum¹⁰. Vogel et al. reported a 12-month incidence rate of 23.7% (n = 350) among a cohort of women in New Zealand⁸. The lower incidence of puerperal mastitis among lactating mothers in our study could be due to two reasons: firstly, exclusive breastfeeding is universal in our study area and secondly, we collected data only up to six weeks after delivery.

The incidence of lactational mastitis was lower among the vaginal delivery patients (4.8%) compared to patients who underwent cesarean section (5.8%) in our study. Prelacteal feeding and cesarean section were associated with a higher likelihood of mastitis⁸. A negative association between Caesarean delivery and breastfeeding exists because postoperative care routines delay the onset of lactation, disrupt mother infant interaction and inhibit infant suckling¹¹. Mothers who had cesarean section were more likely to have problems related to breastfeeding, including nipple fissure, in comparison to women who had vaginal delivery^{12, 13}. However, there were no relationships observed between nipple injuries with the type of childbirth in a cohort study conducted in Australia with 340 primiparous women¹⁴.

In our study, mean age was 27 years of age (Range 18 to 36 years). The majority of the women were between 24-30 years (55%). Maternal age is not directly linked to lactational mastitis; however, there is evidence that young women find it more difficult to breastfeed due to insecurity and inexperience. This can lead to difficulty in breastfeeding, culminating in breast problems and consequently in early weaning¹⁵.

In our study, higher percentage of lactational mastitis was found in primiparous women (57%) compared to multiparous women (43%). Our study is similar to Viduedo AFS et al, reported higher percentage (64%) of severe lactational mastitis among young primiparous women¹⁶. The binational study also reported mastitis more common in primiparous women¹⁷. Women who never breastfed may be more anxious and, as a result, this may interfere with the breastfeeding process¹⁶.

Out of 160 mastitis patients, 72 women were anemic. According to ICMR classification, 13 patients had mild anemia, 40 patients had moderate anemia and 19 patients had severe anemia. Anemic women might be more vulnerable to infection.

Risk factors that have been suggested to be strongly associated to mastitis include cracked or sore nipples, use of ointments, inappropriate breastfeeding practices, mastitis with a previous child, and peripartum antibiotic therapy^{1,7}. The present study is similar to various other studies, cracked nipple or sore nipple were associated with mastitis under the hypothesis that it provides a portal

of entry for microorganisms^{1,2,4,9,18}.

Regarding the amount of milk produced, milk over or undersupply versus normal supply that mastitis may arise from a higher milk supply because of the risk of milk stasis, if the infant delays or misses feeds, this situation may provide good conditions for bacterial overgrowth. On the other hand, low milk supply could give to the mother a false perception of low milk production when, actually, only secretion is compromised due to the formation of thick bacterial biofilms inside the milk ducts^{1,2,18,19}.

The incorrect handling of the infant to the mother's breast and the inadequate positioning between mother and child were associated to nipple trauma. Prevention of nipple damage is likely to reduce the incidence of infectious mastitis. New mothers need good advice about optimal attachment of the baby to the breast and access to skilled help in the early postpartum days and weeks^{13, 20, 21}.

Breast milk culture report of mastitis yielded growth of *Staphylococcus aureus* in 120 (75%) patients and MRSA in 40(25%) patients. Seven patients with mastitis developed breast abscess, the incidence was 4.4% (7/160) as complication. The literature shows that the incidence of breast abscess in lactating mothers varies between 3% and 11%^{6, 18, 24}. In one of the patients with breast abscess, one liter of pus was drained with radial incision made over the breast. Breast Abscess milk culture yielded growth of MRSA and *Staphylococcus aureus*. The most common bacterium found in breast abscess secretion culture was *Staphylococcus aureus*, which coincides with our findings^{15, 23-28}. Various other studies also reported MRSA in breast abscess secretion culture which is similar to our study^{25, 27}. A lactational breast abscess is usually bacterial in etiology and can be effectively managed with oral antibiotics. All patients in our study received antibiotics based on their sensitivity pattern and were discharged in 3 to 4 days. In addition to antibiotics, management of breast abscess included incision and drainage, symptomatic treatment with analgesia and antipyretics, reassurance, assessment of the infant's breastfeeding technique, education, emotional support, and support for continuous breastfeeding.

Conclusion

It is recommended to educate the mothers and the family on the exclusive breast feeding practice and its

benefits, demand feeding, optimal positioning and attachment of the baby, to provide adequate emotional support and access to skilled help in the early postpartum period. Continued breastfeeding should be encouraged in the presence of mastitis which generally does not pose a risk to the infant. Breast abscess is the most common complication of mastitis. It can be prevented by early treatment of mastitis and continued breastfeeding.

Conflict of interest: None. **Disclaimer:** Nil.

References

1. Amir LH, Forster DA, Lumley J and McLachlan H. A descriptive study of mastitis in Australian breastfeeding women: incidence and determinants. BMC Public Health. 2007; 7: 62.
2. Cullinane M, Amir L, Donath SM. Determinants of mastitis in women in the CASTLE study: a cohort study. BMC Family Practice. 2015; 16: 181.
3. Konar H. Abnormalities of the Puerperium. In: Konar H, editor. DC Dutta's Textbook of Obstetrics. 8th ed. New Delhi: Jaypee; 2015: p500-13.
4. Amir LH, Garland SM, Lumley J. A case-control study of mastitis: nasal carriage of *Staphylococcus aureus*. BMC Family Practice. 2006; 7: 57.
5. Iyengar K. Early Postpartum Maternal Morbidity among Rural Women of Rajasthan, India: A Community-based Study. J Health Popul Nutr. 2012 Jun; 30(2): 213-25.
6. Cusack L, Brennan M. Lactational mastitis and breast abscess: diagnosis and management in general practice. Aust Fam Physician. 2011; 40(12): 976.
7. Spencer JP. Management of mastitis in breastfeeding women. Am Fam Physician. 2008 sep 15; 78(6): 727-31.
8. Khanal V, Scott JA, Lee AH, Binns CW. Incidence of Mastitis in the Neonatal Period in a Traditional Breastfeeding Society: Results of a Cohort Study. Breastfeeding Medicine. 2015; 10(10): 481-7.
9. Tang L, Lee AH, Qiu L, Binns CW. Mastitis in Chinese breastfeeding mothers: a prospective cohort study. Breastfeed Med. 2014 Jan-Feb; 9(1): 35-8.
10. Scott JA, Robertson M, Fitzpatrick J, Knight C, Mulholland S. Occurrence of lactational mastitis and medical management: A prospective cohort study in Glasgow. Int Breastfeed J. 2008; 3: 21.
11. Prior E, Santhakumaran S, Gale C, Philipps LH, Modi N, Hyde MJ. Breastfeeding after cesarean delivery: a systematic

review and meta-analysis of world literature. *Am J Clin Nutr.* 2012; 95: 1113-35.

12. Boskabadi H, Ramazan-zadeh M, Zakerihamidi M, Rezagholizade OF. Risk factors of breast problems in mothers and its effects on newborns. *Iran Red Crescent Med J.* 2014; 16 (6): e8582.

13. Silva DJ, de Oliveira VT, Oliveira VG. Factors associated to nipple trauma in lactation period: a systematic review. *Rev Bras Saude Mater Infant.* 2017 Mar;17(1): 27-42.

14. Buck ML, Amir LH, Cullinane M, Donath SM. Nipple pain, damage, and vasospasm in the first 8 weeks postpartum. *Breastfeed Med.* 2014; 9 (2): 56-62.

15. Kinlay JR, O'Connell DL, Kinlay S. Risk factors for mastitis in breastfeeding women: results of a prospective cohort study. *Aust N Z J Public Health.* 2001; 25: 115-20.

16. Viduedo AFS, Leite JRC, Monteiro JCS, Reis MCG, Gomes-Sponholz FA. Severe lactational mastitis: particularities from admission. *Rev Bras Enferm.* 2015; 68(6): 806-11.

17. Iatrakis G, Zervoudis S, Ceausu I, Peitsidis P, Tomara I, Bakalianou K, Hudita D. Clinical features and treatment of lactational mastitis: the experience from a binational study. *Clin Exp Obstet Gynecol.* 2013; 40(2): 275-6.

18. Mediano P, Fernández L, Rodríguez JM, Marín M. Case control study of risk factors for infectious mastitis in Spanish breastfeeding women. *BMC Pregnancy and Childbirth.* 2014; 14: 195.

19. Delgado S, Arroyo R, Jiménez E, Marín ML, del Campo R, Fernández L, Rodríguez JM. *Staphylococcus epidermidis* strains isolated from breast milk of women suffering infectious mastitis: potential virulence traits and resistance to antibiotics. *BMC Microbiol.* 2009; 9: 82.

20. Kronborg H, Vaeth M. How Are Effective Breastfeeding Technique and Pacifier Use Related to Breastfeeding Problems and Breastfeeding Duration? *Birth.* 2009; 36: 34-42.

21. Goyal RC, Banginwar AS, Ziyof, Toweir AA. Breastfeeding practices: Positioning, attachment (latch-on)

and effective suckling-A hospital-based study in Libya. *J Fam Comm Med.* 2011; 18 (2): 74-9.

22. Bertrand H, Rosenblood LK. Stripping out pus in lactational mastitis: a means of preventing breast abscess. *Can Med Assoc J.* 1991; 145: 299-306.

23. Kvist LJ, Larsson BW, Hall-Lord ML, Steen A, Schalen C. The role of bacteria in lactational mastitis and some considerations of the use of antibiotic treatment. *Int Breastfeed J.* 2008; 3: 6.

24. Amir LH, Forster D, McLachlan H, Lumley J. Incidence of breast abscess in lactating women: report from an Australian cohort. *BJOG.* 2004; 111(12): 1378-81.

25. Ramakrishnan R, Trichur RV, Murugesan S, Cattamanchi S. Analysis of the microbial flora in breast abscess: a retrospective cohort study conducted in the emergency department. *Int Surg J.* 2017; 4: 2143-7.

26. Sandhu GS, Gill HS, Sandhu GK, Gill GP, Gill AK. Bacteriology in breast abscesses. *Sch J App Med Sci.* 2014; 2(4E): 1469-72.

27. Chick JFB, Chauhan NR, Polen LA. Emergency management of puerperal breast abscess. *Intern Emerg Med.* 2012; 7(2): S167-S168.

28. Kataria K, Srivastava A, Dhar A. Management of Lactational Mastitis and Breast Abscesses: Review of Current Knowledge and practice. *Indian J Surg.* 2013 Nov-Dec; 75(6):115-20.

Ruma Dutta¹, Ravikanth Onkarappa Gowder²

¹Senior Resident, Dept. of Obstetrics and Gynaecology, Aditya Diagnostics and Hospital, Dibrugarh, Assam, India; ²Associate Professor, Dept. of Obstetrics and Gynaecology, K.V.G Medical College and Hospital, Sullia, Karnataka, India.