

Obstetric outcome of elevated total serum bile acid levels in women with intrahepatic cholestasis of pregnancy

Vineeta Gupta, Alvina Rehman, Shweta Nimonkar, Priyanka Chaudhari, Namrata Saxena

Corresponding author: Dr. Vineeta Gupta, Prof & Head, Department of Obstetrics and Gynaecology, SGRIHM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India; Email : vineetahims@yahoo.co.in

Distributed under Attribution-Non Commercial – Share Alike 4.0 International (CC BY-NC-SA 4.0)

ABSTRACT

Background: Intrahepatic cholestasis of pregnancy (IHCP) is a hepatic disorder associated specifically with pregnancy. It is found to affect both mother and fetus. It requires detailed clinical and biochemical evaluation for optimal fetomaternal outcome. **Objectives:** To study the demographic profile, clinical features and obstetric outcome in patients with intrahepatic cholestasis of pregnancy and to assess correlation between serum bile acid levels and perinatal outcome. **Material and methods:** This was a prospective study conducted on 150 pregnant women with IHCP who delivered at our tertiary care centre, Shri Mahant Indires Hospital, Dehradun from September 2018 – June 2020. The relationship between serum bile acid levels and fetal complications was analysed with binary logistic regression method. **Results:** The incidence of IHCP at our centre was found to be 2.82%. We found that 59.94% patients had bile acid levels between 10-40 $\mu\text{mol/L}$, 36.6% patients bile acid levels between 41-100 $\mu\text{mol/L}$ and 3.33% patients had bile acid levels $>100 \mu\text{mol/L}$. The correlation between serum bile acid levels with onset of labour and mode of delivery was not statistically significant (p-value 0.16 and 0.556 respectively). The correlation between serum bile acid levels with meconium stained liquor, NICU admission and prematurity was found to be statistically significant with p-value of <0.001 , 0.007 and 0.018 respectively. **Conclusion:** Symptomatic women should be subjected to estimation of serum bile acid levels so that appropriate treatment and timely intervention can be done to optimize obstetric outcome.

Keywords: Intrahepatic cholestasis of pregnancy, pruritis, serum bile acids, meconium stained liquor, ursodeoxycholic acid.

Intrahepatic cholestasis (IHCP) of pregnancy is a hepatic disorder in the late second and early third trimester of pregnancy. It is characterized by maternal pruritis in absence of skin rash with increased serum bile acids and elevated serum aminotransferases. Overall incidence of IHCP is between 0.02% to 2.4%¹. The etiopathophysiology of IHCP is still not clearly understood. The clinical symptoms and biochemical abnormality resolves after delivery. IHCP is associated with increased risk of adverse obstetrical outcomes, which includes stillbirth, meconium passage, respiratory distress syndrome and fetal asphyxia¹. It is found that elevated serum bile acids levels in the fetal circulation

plays significant role in fetal complications. Patients are classified as mild (S. Bile acids 10 – 40) or severe (S. Bile acids >40). Various studies have suggested that the risk of fetal complications is directly proportional to the increasing levels of serum bile acids^{2,3}.

In view of limited studies on IHCP and its correlation with serum bile acids, the present study was conducted to study the clinical presentation and obstetric outcome in patients with IHCP. In our study, the correlation between serum bile acids and perinatal outcome was also assessed.

Materials and methods

This was a prospective study of maternal and perinatal

Received: 5th September 2021, Peer review completed: 16th December 2021, Accepted: 30th December 2021.

Gupta V, Rehman A, Nimonkar S, Chaudhari P, Saxena N. Obstetric outcome of elevated total serum bile acid levels in women with intrahepatic cholestasis of pregnancy. The New Indian Journal of OBGYN. 2023; 9(2): 302 - 7.

consequences of elevated bile acid levels in pregnant women presenting with intrahepatic cholestasis of pregnancy, which was carried out from September 2018 to June 2020 in Department of Obstetrics & Gynaecology, at Shri Mahant Indires Hospital, Dehradun after clearance from the institutional ethics committee. A total of 150 patients with intrahepatic cholestasis of pregnancy were included. Diagnosis was based on presence of pruritus, elevated total serum bile acid levels and serum transaminases. Following patients were excluded from the study:

- a. Patients with one or more medical diseases such as cardiologic, oncologic, hematologic, acute and chronic kidney diseases.
- b. Patients having other liver disorders like viral hepatitis A, B, C, D, E.
- c. Patients having cholelithiasis, bile duct lithiasis, cholangitis, other abnormalities of liver and biliary tract, alcohol addiction, HIV infection, substance abuse.
- d. Patients with diseases of skin with itching and rashes.

After informed consent, a thorough history was taken. Complete general and obstetrical examination was carried out on the patients. Besides routine investigations, the patients were also subjected to following biochemical tests-

- i) Aspartate aminotransferase (AST)
- ii) Alanine aminotransferase (ALT)
- iii) Alkaline phosphatase
- iv) Total serum bile acids
- v) Total serum bilirubin

Aminotransferases were analysed using UV with PSP (dry chemistry) method, alkaline phosphatase was analysed using PNPP, AMP (dry chemistry) method, total serum bile acid levels by enzyme assay calorimetry method, total serum bilirubin level by diphylline (dry chemistry) method.

Based on investigations and clinical assessment, patients were treated with ursodeoxycholic acid, antihistamines and topical emollients. Repeat samples were advised for patients after 7-14 days depending on symptoms and severity of disease. Decision regarding further management was based on worsening of symptoms and serial estimation of serum bile acid levels. Patients were monitored by regular antenatal visits with daily maternal recording of fetal movements, serial sonography for fetal well being, twice weekly fetal cardiotocography (CTG) from 34 weeks of gestation. Steroid coverage was given for fetal lung maturity for suspected preterm deliveries before 34weeks of gestation and vitamin K injection was given before delivery. Given the increased

risk of stillbirth in the setting of IHCP, pregnancies were actively managed by timely delivery, through induction of labor as recommended at around 37 weeks of gestation or earlier to balance the risk of iatrogenic preterm delivery against the risk of fetal mortality. However, these decisions were individualized with careful patient counselling. Obstetric outcome was recorded in every case in terms of maternal and fetal outcome.

Results thus obtained were tabulated and analyzed. The relationship between serum bile acid levels and transaminases level, and that between serum bile acid levels and fetal complication was analysed with binary logistic regression method. Standard statistical methods were applied, according to Chi-square method, p-value< 0.05 was considered to be significant.

Results

During the study period, there were 5316 deliveries conducted at our tertiary care hospital. Out of these, 150 patients with intrahepatic cholestasis of pregnancy were included in this study. The overall incidence of intrahepatic cholestasis of pregnancy recorded at our hospital was found to be 2.82%. Table 1 shows the demographic profile and clinical presentation of the cases. All the patients presented with complaints of pruritus/itching either on whole body or

Table 1: Demographic profile and clinical features of cases		
Categories	Number	Percentage (%)
Age (yrs)		
≤ 25	40	26.66
26-30	77	51.33
31-35	28	18.66
>35	5	3.33
Socio-economic status (Modified Kuppuswamy scale)		
I (Upper)	20	13.33
II (Upper middle)	14	9.33
III (Lower middle)	89	59.33
IV (Upper lower)	22	14.67
V (Lower)	5	3.33
Locality		
Rural	53	35.33
Urban	97	64.67
Gravida		
G1	76	50.66
G2	36	24
G3	28	18.66
G4	5	3.33
G5	5	3.33
Clinical features		
Pruritus	150	100
Excoriation/Rash	68	45.33
Jaundice	3	2

palms and soles. It was more at night time which resulted in insomnia in many patients. Excoriation or rashes were observed in 68 (45.33%) patients whereas clinical jaundice was seen only in 3 (2%) patients.

Table 2: Showing initial and serial serum bile acid levels and LFTs

Initial S. bile acids (µmol/L)	Number	Percentage (%)
10-40	90	59.94
41-100	55	36.66
>100	5	3.33
Total	150	100
Serial S. bile acids (µmol/L)		
10-40	16	30.77
41-100	34	65.38
>100	2	3.85
Total	52	100
Liver function tests		
	Reference value	Mean
S. bilirubin [mg/dl]	0.2-1.3	0.625 ± 0.325
AST (SGOT) [U/L]	17-59	108.46 ± 41.08
ALT (SGPT) [U/L]	21-72	110.20 ± 38.40
ALP [U/L]	38-126	350.2 ± 175.34

Serum bile acid levels were estimated in all the patients. As seen in table 2, ninety (59.94%) patients had initial bile acid levels between 10-40 µmol/L, 55 (36.6%) patients between 41-100 µmol/L and only 5 (3.33%) patients had levels >100 µmol/L. Serial estimation of serum bile acid levels were advised for patients with worsening symptoms, especially with bile acid levels > 40 µmol/L. Out of 150 patients, only 52 patients had repeat estimation of serum bile acid levels. Ninety-eight patients did not have repeat test either because of high cost of test or due to non-compliance or their symptoms improved. Repeat S. bile acid levels were found to be increasing in 33 patients, decreasing in 14 and remained same in 5 patients. In patients with increasing levels, termination of pregnancy was advised after counselling. Mode of termination was decided based on individual patient's profile, either by induction of labour or caesarean section for other obstetric indications. Patients with decreasing or same S. bile acid levels were monitored and followed up with regular antenatal visits, biochemical studies and CTG monitoring after explaining prognosis of the condition. So, serial bile acid level estimation helped in deciding management of these patients.

Liver function tests were monitored for all the patients. Serum bilirubin was usually within reference value for the patients, with mild increase in AST and ALT (table 2). Alkaline phosphatase was increased in majority of the patients. Mean S. bilirubin, AST, ALT, ALP values were 0.625 ± 0.325, 108.46 ± 41.08, 110.20 ± 38.40 and 350.2 ± 175.34 respectively. In 87 (58%) patients, LFTs were within

normal limits, whereas they were deranged in 63 (42%) patients. After starting pharmacotherapy, LFTs were repeated in 63 patients in whom the values were abnormal initially. In 54 (85.71%) patients values improved and deteriorated in 11 (14.29%) patients. Symptoms improved in 133 (88.67%) patients after starting pharmacotherapy. Eleven (7.33%) patients showed no signs of improvement and worsening of symptoms was seen only in 6 (4%) patients.

Table 3 shows correlation between S. bile acid levels with onset of labour and mode of delivery. Out of 150 deliveries, 85 were vaginal deliveries, 16 were elective caesarean sections and 49 were emergency caesarean sections. Out of 85 vaginal deliveries, 6 were instrumental vaginal deliveries. Indications of intervention have been mentioned in table 4. In majority patients, induction of labour was done for IHCP (82.19%), followed by PROM in 9.59% patients. Elective caesarean section was done in 16 patients, most common indication being previous 1 caesarean delivery with refusal for trial of labour in present pregnancy (43.75%). Emergency caesarean section was done in 49 patients for various indications, most common

Table 3: Correlation of S. bile acid levels with onset of labour and mode of delivery

S. bile acid (µmol/L)	Total	Onset of labour and mode of delivery					
		Vaginal		Caesarean		Emergency No (%)	
		Spontaneous Normal No (%)	Instrumental No (%)	Induced Normal No (%)	Instrumental No (%)		Elective No (%)
0 - 40	90	21(23.33%)	2(2.22%)	27(30%)	3(3.33%)	6(6.7%)	31(34.44%)
41 - 100	55	10(18.2%)	0(0%)	18(32.72%)	1(1.82%)	9(16.36%)	17(30.90%)
>100	5	2(40%)	0(0%)	1(20%)	0(0%)	1(20%)	1(20%)
Total	150	33(22%)	2(1.33%)	46(30.67%)	4(2.67%)	16(10.67%)	49(32.67%)

indication being fetal distress in 25 (51.02%) patients. Mean hospital stay of patients (in days) with S. bile acid levels 0-40, 41-100, >100 µmol/L was 5.74 ± 5.43, 5.49 ± 3.24, 5.00 ± 2.55 respectively. The correlation between S. bile acid levels with mean hospital stay of mother was statistically insignificant (p-value 0.908). There were 141 uneventful deliveries without any complication. One patient needed ICU care post-operatively and 8 patients had PPH which was medically managed. No surgical intervention was required in these 8 patients. The correlation between S. bile acid levels with maternal complications during delivery was statistically insignificant (p-value 0.422).

There were 158 neonates born to 150 mothers as there were 8 twin pregnancies. There were 2 intra-uterine fetal deaths. Out of 156 live births, 102 (65.38%) neonates were normal and did not have any complications, while 54

Table 4: Indications for induction of labor and caesarean delivery

Indications	Number (%)
Induction of labour	
IHCP	60 (82.19%)
PROM	7 (9.59%)
Hypertensive disorders	2 (2.74%)
Postdatism	1 (1.37%)
GDM	2 (2.74%)
IUD	1 (1.37%)
Elective caesarean section	
Previous 1 CS with refusal of TOLAC	7 (43.75%)
Caesarean delivery on maternal request	6 (37.5%)
Transverse lie	1 (6.25%)
Twin gestation with 1 st twin breech	2 (12.5%)
Emergency caesarean section	
Fetal distress	25 (51.02%)
Previous 1 CS in labour with refusal of TOLAC	4 (8.16%)
Previous 2 CS in labour	2 (4.08%)
Placenta previa with APH	2 (4.08%)
Primi breech in labour	4 (8.16%)
Non progress of labour	3 (6.12%)
Previous CS with twin in labour	1 (2.04%)
Transverse lie in labour	2 (4.08%)
Second stage arrest of labour	2 (4.08%)
Previous CS with scar dehiscence	1 (2.04%)
Twin gestation with 1 st twin breech in labour	2 (4.08%)
Previous CS with breech in labour	1 (2.04%)
Twin gestation with cord prolapse	1 (2.04%)

(34.62%) neonates had one or more problems as seen in table 5. Meconium staining of liquor was seen in 37 (24.67%), birth asphyxia in 13 (8.3%) neonates, 54 (34.6%) required NICU admission and 51 (34%) were born prematurely (<37 weeks). The correlation between S. bile acid levels with meconium stained liquor, NICU admission and prematurity was found to be statistically significant with p-value of <0.001, 0.007 and 0.018 respectively. Correlation of S. bile

Table 5: Correlation of S. bile acid levels with fetal outcome

S. bile acid (µmol/L)	0 - 40	41 - 100	>100	Total	P value
Meconium stained liquor [Number (%)]	13 (35.14%)	20 (54.05%)	4 (10.81%)	37 (24.67%)	< 0.001
Apgar Score					
At 1 min	7.12 ± 1.03	7.11 ± 1.15	7.20 ± 0.84	-	0.983
At 5 min	8.18 ± 1.06	8.20 ± 1.25	8.20 ± 0.84	-	0.993
Birth asphyxia [Number (%)]	7 (53.84%)	5 (38.46%)	1 (7.7%)	13 (8.3%)	0.830
NICU admission [Number (%)]	24 (44.44%)	26 (48.15%)	4 (7.41%)	54 (34.6%)	0.007
Prematurity[Number (%)]	23 (45.1%)	24 (47.06%)	4 (7.84%)	51 (34%)	0.018
IUD [Number (%)]	1 (50%)	-	1 (50%)	2 (1.27%)	-

acid levels with mean Apgar score at 1 and 5 minutes and birth asphyxia was found to be insignificant (p-value 0.983, 0.993 and 0.830 respectively). Regarding birth weight of neonates, 110 had birth weight >2500 gm, out of these 17 were preterm and 93 were term babies. Forty eight were low birth weight babies (<2500 gm), out of these 32 were preterm and 16 were term babies. Low birth weight for these 16 term babies was attributed to gestational hypertension in 3, uteroplacental insufficiency in 5, placenta previa in 2, multiple gestation in 1, polyhydramnios in 1 and idiopathic in 4 babies. No baby weighed <1000 gm.

Discussion

In our study, the incidence was of 2.82%. The incidence was found to be 2.73% by Pegu B et al⁴, 2.4% by Sharma et al⁵ and 2.59% by Alakananda et al⁶. The mean age of patients in our study was 27.85 years. It was comparable to a recent study by Joseph B et al⁷. Majority of the patients belonged to lower middle class group (59.3%) according to modified Kuppaswamy scale (2017) and resided in urban locality (64.67%). Other authors have not taken into account the socio-economic scale and locality, so we could not compare these demographic data. In our study, 50.66% (n = 76) were primipara and 49.34% (n = 74) were multiparous. In a study by Joseph B et al⁷, 68.3% were primipara and 31.7% were multiparous.

In our study, pruritus was seen in 100% patients which was their presenting complaint, excoriation marks or rashes were seen in 45.33% patients. Pegu B et al⁴ in their study found that main symptom of IHCP was pruritus (88%) which cause disturbed sleep (35%) and scratch marks (73%). It has been reported that itching may be presented either before or after abnormal liver function test and this may reflect the heterogeneous nature of this condition.

Study conducted by Alakananda et al found serum bile acids and liver enzymes were raised in 100% cases⁶. In our study, 59.94% (n=90) patients had bile acid levels between 10-40 µmol/L, 36.6% (n=55) patients had levels between 41-100 µmol/L and only 3.33% (n=5) patients had levels >100 µmol/L.

In our study 40.66% (n=61) had spontaneous labour, in 47.33% (n=71) patients induction of labour was done and for 16 (10.67%) patients elective caesarean section was done before onset of labour for various indications. Pegu B et al found that most of the patients went into labour spontaneously and some of them needed induction⁴. Some studies have reported good outcomes with a policy of induction of labour at 37 or 38 wk gestation. Many clinicians in the UK have adopted this practice as the IUDs appear to cluster at later gestations².

Rate of instrumental delivery and caesarean delivery is higher in pregnancies affected by IHCP due to increased

incidence of meconium staining of liquor and induction of labour. In our study, 85 patients (56.7%) had vaginal delivery and 65 patients (43.3%) underwent caesarean section. Out of 85 deliveries, 6 (4%) were instrumental deliveries. Maximum number of caesarean sections were done for 49.1% (n=27) patients with serum bile acid levels 41-100 $\mu\text{mol/L}$. Comparable results were seen in a study by Sharma et al⁵, where normal delivery occurred in 50% patients, instrumental delivery in 8.33% patients and caesarean section in 41.66% patients. Pegu B et al⁴ found that normal delivery occurred in 60.29% patients, instrumental delivery in 8.82% and caesarean section in 30.8% patients.

In our study the incidence of PPH was found to be high in women with IHCP i.e. 5.3% (n = 8) which was comparable to Alakananda et al study with 6% cases of PPH⁶. Other studies did not mention about maternal ICU stay. In our study, only 1 (0.67%) patient had to stay in ICU for observation.

Even though IHCP is benign to the mother, it is associated with increased fetal complications. In our study we found association with perinatal complications, including meconium-stained amniotic fluid in 24.6%, NICU admissions in 32.7% and there were 2 (1.3%) unbooked cases who presented with intrauterine fetal death. In comparison, Alakananda et al found meconium-stained amniotic fluid in 29%, NICU admissions in 21% and intrauterine death in 2% cases⁶. The incidence of preterm delivery in our study was 24.66% compared to 23% in study by Alakananda et al⁶. Logistic regression analyses of the data for fetal complications suggested that meconium staining of liquor, green placenta, NICU admissions for neonates increased as levels of bile acid increased. Fetal outcome in different studies by Kenyon AP et al⁸, Glantz A et al⁹, Singh G et al¹⁰, Gupta A et al¹¹, Dang A et al¹², Padmaja M et al¹³ and Alakananda et al⁶ is seen in table 6.

Glantz A et al reported increased adverse fetal outcome if serum bile acid was $>40 \mu\text{mol/l}$ and no increase in adverse outcomes if the maternal fasting serum bile acids were below $40 \mu\text{mol/L}$, leading the authors to suggest there is no increased risk to the fetus with mild IHCP⁹. In our study also, we found that patients having serum bile acid level $>40 \mu\text{mol/l}$ seem to have more fetal complications (p value 0.0007). Hani A et al¹⁴ reported that significant positive correlation was seen between maternal fasting bile acids level and meconium staining of amniotic fluid (P value <0.001) which was found in our study also (p value <0.001).

However, in spite of high incidence of meconium staining, Apgar score is usually normal. It shows that meconium staining of amniotic fluid seen in IHCP is not due to fetal distress; instead it is because of bile acids as bile acids are suggested to increase colonic motility and meconium. Mean birth weight was $2.6 \pm 0.4 \text{ kg}$ which is appropriate for gestational age. It indicates that birth weight is not affected by IHCP. Similar to our study, other studies also did not find any significant maternal adverse outcome whereas association with poor perinatal outcome was significant correlating with increasing serum bile acid levels.

Conclusion

The incidence of cholestasis of pregnancy is high in the Indian population. Serum bile acid levels are increased in cholestasis of pregnancy. Poor perinatal outcome is associated with derangements in serum bile acid levels and liver function tests. Proper monitoring during antenatal period and termination of pregnancy at term improves the perinatal outcome. In our study, significant statistical correlation has been found between serum bile acid levels and prematurity, meconium staining of liquor and NICU admission of neonates. Though, we did not find any significant correlation between serum bile acid levels and Apgar score, birth asphyxia and average NICU stay. Serial bile acid levels estimation helped in deciding management of these patients. We recommend that symptomatic patients should be subjected to estimation of serum bile acid levels and liver transaminases, so that appropriate treatment can be provided and timely intervention can be done to optimize obstetric outcome.

Conflict of interest: None. **Disclaimer:** Nil.

References

1. Chacko KR, Wolkoff AW. Intrahepatic cholestasis of pregnancy: new diagnostic insights. *Ann Hepatol.* 2017; 16 (2): 176-78.
2. Geenes V, Williamson C. Liver disease in pregnancy. *Best Pract Res Clin Obstet Gynecol.* 2015; 29: 612-24.
3. Brouwers L, Koster MP, Page-Christiaens GC, Kemperman H, Boon J, Evers IM, et al. Intrahepatic cholestasis of pregnancy: maternal and fetal outcomes associated with elevated bile acid levels. *Am J Obstet Gynecol.* 2015; 212: 100. e1-e7.
4. Pegu B, Manju M, Anita Y, Sahoo PSK. Cholestasis of pregnancy: a prospective analysis from a South

- Andaman teaching hospital. *Int J Reprod Contracept Obstet Gynecol*. 2019; 8:1895-8.
5. Sharma N, Panda S, Santa Singh A. Obstetric outcome during an era of active management for obstetrics cholestasis. *J Obstet Gynecol India*. 2015; 66(1): 38-41.
 6. Alakananda, Bhattacharya A, Kavita. Feto-maternal Outcome in intrahepatic cholestasis of pregnancy. *Sch J App Med Sci*. 2016; 4(10D): 3837-41.
 7. Joseph B, et al. Obstetric Outcome in Women with Intrahepatic Cholestasis: A 3-year Study in a Tertiary Care Hospital in Bengaluru. *J South Asian Feder Obst Gynae*. 2019; 11(2): 103-6.
 8. Kenyon AP, Piercy CN, Girling J, Williamson C, Tribe RM, Shennan AH. Obstetric cholestasis outcome with active management: a series of 70 cases. *BJOG*. 2002; 109: 282-88.
 9. Glantz A, Marschall HU, Mattsson LA. Intrahepatic cholestasis of pregnancy: relationships between bile acid levels and fetal complication rates. *Hepatology*. 2004; 40: 467-74.
 10. Singh G, Sidhu K. Cholestasis of Pregnancy: A Prospective Study. *Med J Armed Forces India*. 2008; 64(4): 343-45.
 11. Gupta A, Tania K, Yudhishtervir G, Jyoti H. Cholestasis of pregnancy. *J Obstet Gynecol India*. 2009; 59: 320.
 12. Dang A, Agarwal N, Bathla S, Sharma N, Balani S. Prevalence of liver disease in pregnancy and its outcome with emphasis on obstetric cholestasis: An Indian scenario. *The Journal of Obstetrics and Gynecology of India*. 2010 Oct 1; 60(5): 413-8.
 13. Padmaja M, Bhaskar P, Kumar GJ, Seetha R, Mahasweta C. A study of obstetric cholestasis. *J Obstet Gynecol India*. 2010; 60(3): 225-31.
 14. Al Shobaili HA, Hamed HO, Al Robaee A, Alzolibani AA, Amin AF, Ahmad SR. Obstetrical and fetal outcomes of a new management strategy in patients with intra-hepatic cholestasis of pregnancy. *Archives of Gynecology and Obstetrics*. 2011 Jun 1; 283(6): 1219-25.
-
- Vineeta Gupta¹, Alvina Rehman², Shweta Nimonkar³, Priyanka Chaudhari⁴, Namrata Saxena⁵**
¹ Prof & Head, Department of Obstetrics and Gynaecology, SGRRIM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India; ² Junior Resident, Department of Obstetrics and Gynaecology, SGRRIM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India; ³ Associate Professor, Department of Obstetrics and Gynaecology, SGRRIM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India; ⁴ Professor, Department of Obstetrics and Gynaecology, SGRRIM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India; ⁵ Associate Professor, Department of Obstetrics and Gynaecology, SGRRIM&S & Shri Mahant Indires Hospital, Patel Nagar, Dehradun, Uttarakhand, India.