

Prevalence of meconium stained liquor in postdated pregnancy in a tertiary care hospital in India

Shatakshi Kulshreshtha, Sujata Siwatch

Corresponding author: Dr Sujata Siwatch, Associate Professor, Department of Obstetrics and Gynecology, PGIMER Chandigarh, India; Email: sujatasiw@yahoo.com

Distributed under Attribution-Non Commercial – Share Alike 4.0 International (CC BY-NC-SA 4.0)

ABSTRACT

Objective: To determine the prevalence of meconium stained liquor in women at and beyond expected date of delivery in Indian population. **Methods:** A retrospective analysis of all women delivered at more than 40 weeks from January 2017 to December 2018 in PGIMER was done. Data was collected from labour room registries. **Results:** A total of 11,675 women delivered in the labor room at our institute during the year 2017-2018. Of these, 717(6.14%) were more than 40 weeks. Meconium stained liquor (MSL) was present in 107 (14.92%). In our study, 86.9% of women with MSL underwent a cesarean section while 7/107 (6.5%) had instrumental delivery. **Conclusion:** Prevalence of MSL increases with increasing gestational age after 40 weeks. Most of the pregnancies with MSL eventually delivered via cesarean section.

Keywords: Meconium, postdated pregnancy, meconium aspiration syndrome, induction, fetal distress.

Meconium stained amniotic fluid is important in influencing obstetrician's decision as it has long been thought to be an indicator of fetal distress. Meconium stained liquor can increase neonatal respiratory morbidity by causing meconium aspiration syndrome (MAS). This syndrome develops in around 1% of the cases¹. Meconium stained amniotic fluid is also associated with other complications like sepsis, neurological impairment, convulsions and prolonged NICU admissions². More advanced gestational age presents a greater risk of meconium stained liquor (MSL).

MAS is a major cause of perinatal morbidity and mortality in postdated pregnancies. It has been thought that intrauterine hypoxia may result in anal sphincter relaxation leading to passage of meconium. Physiologically, fetal maturity of parasympathetic nervous system leads to sphincter relaxation and passage of meconium. Caughey AB et al³ demonstrated a level of meconium in amniotic fluid six times greater among women at 42 weeks or more, compared with those at 37 weeks. In a study done by

Caughey et al⁴ there is increased risk of cesarean section and meconium stained liquor beyond 41 weeks. Hence, prolonged pregnancy is one of the common reasons for induction of labour.

Postdated pregnancy is the one which has crossed expected date of delivery. Post term pregnancy is the one that has completed or gone beyond 42 weeks of gestation⁵. The prevalence of late term and post term births in United States is around 7%⁶. A number of observational studies have identified risk factors for prolonged pregnancy to include primigravidity, prior post-term pregnancy, male fetus, obesity and genetic predisposition⁷. American College of Obstetrics and Gynecology recommends that, women beyond 41 weeks should be offered induction of labour and beyond 42 weeks should be considered for induction⁵. S-1 guidelines by Weiss et al in Germany recommend offering pregnant women the option of inducing labor from 41 + 0 gestation and state that induction of labor must be recommended by 41 + 3 gestation at the latest⁸.

There is evidence of ethnic variations in meconium

Received: 5th June 2021, Peer review completed: 10th July 2021, Accepted: 9th October 2021.

Kulshreshtha S, Siwatch S. Prevalence of meconium stained liquor in postdated pregnancy in a tertiary care hospital in India. The New Indian Journal of OBGYN. 2023; 10(1): 67 -70.

stained amniotic fluid in term pregnancies. This leads to different rates of stillbirth and perinatal morbidity as the pregnancy approaches late term and beyond. Independent predictors of meconium stained amniotic fluid includes being Black (odds ratio 8.4, 95% CI 2.4 - 28.8) or South Asian (OR 3.3, 95% CI 1.3-8.3) ⁹. Fetuses of Black and Indian races can pass meconium early and consequently can cause fetal distress. This urged us to hypothesize that in Asians and Blacks, fetus attains maturity earlier. The significance is that physicians should increase antenatal surveillance beyond 40 weeks. With this background we aim to find out the prevalence of MSL in Indian population beyond 40 weeks.

Objective: To determine the prevalence of meconium stained liquor in women at and beyond expected date of delivery in Indian population.

Materials and methods

A retrospective analysis of all women delivered at more than 40 weeks from January 2017 to December 2018 in PGIMER, a tertiary care hospital in Northern India was done. All those women who were sure of her last menstrual period (LMP) and had first trimester dating ultrasound scan were included. The cases where LMP was not known and no first trimester USG was available to ascertain the gestational age were excluded. Data were collected from labour room registries.

Other variables associated with the presence of meconium in amniotic fluid like period of gestation at and beyond 40 weeks, induced/spontaneous labor, complications in pregnancy (hypothyroidism, hypertensive disorders, abruption, intrahepatic cholestasis of pregnancy, intrauterine growth retardation), major congenital malformations, mode of delivery and neonatal outcome were also evaluated.

Results

A total of 11,675 women delivered in the labor room at our institute during the year 2017-2018. Of these, 717(6.14%) were more than 40 weeks. The distribution of gestational age of these 717 pregnancies at or beyond 40 weeks is shown in table 1. Labor was induced in 307(42.8%) of these women with gestational age of 40 weeks. The mode of delivery was cesarean section in 289/707(40.30%) of all women. One had peripartum hysterectomy for uterine rupture. Meconium stained liquor was present in 107 (14.92%).

Forty four (41.1%) women who subsequently had MSL were induced either for postdated pregnancy or for other obstetrical indication. Factors like intrauterine growth

restriction, abruption, hypothyroidism and cholestasis that are associated with MSL were present in 40/107(37.3%) while in 67/107 (62.61%) they were not. In our study, 86.9% of women with MSL underwent a cesarean section while 7/107 (6.5%) had instrumental delivery. Congenital malformations were present in 15.9% of all babies delivered, 5.29% being major malformations. Out of 717 deliveries, 17 ended up in stillbirth (2.37%). Out of the 17 stillborn, 47% had major congenital malformation, 17.64% had obstetric cause associated, 5.80% presented with MSL (the probable cause was fetal distress) and 29.41 % were intrauterine deaths on admission. Amongst all the babies born with MSL, only one (0.93%) baby had one minute Apgar score < 6.

Table 1: Distribution of gestational age

Gestational age in weeks (+days)	Pregnancies in number	Pregnancies with MSL	Prevalence of MSL (%)
40 - 40+3	550	78	14.18
40 + 4 to 40 + 6	112	16	14.28
≥ 41	52	12	23.07
≥ 42	10	3	30
≥ 43	3	1	33.3

Discussion

Prevalence of meconium stained amniotic fluid in postdated pregnancies in our institute is 14.92 %. According to a study done by Narasimhaiah A et al in a hospital in Delhi, the incidence of MSL in postdated pregnancies was 11.73% ¹⁰ while Addisiu et al ¹¹ reported a prevalence of MSL of 17.8% in term pregnancies. According to Hirsch et al ¹², the overall prevalence of meconium stained amniotic fluid was 12% in pregnancies at and beyond term. A positive relation was observed between the rates of meconium staining and advancing gestational age. This reinforces the theory that meconium staining of liquor is a physiological event and is linked to maturity of fetal gastrointestinal tract. But still, presence of meconium is a risk factor for respiratory morbidity. The highest odds for respiratory morbidity were found to be in late term gestation when pregnancies with clear liquor were compared with meconium stained liquor. They also found that the effect of meconium stained liquor and increased respiratory morbidity is most significant at 41 weeks of gestation.

In our study, most of the babies of women with MSL had Apgar score of >7 at 1minute and 5 minutes. Although the prevalence of meconium stained liquor increased as the gestation advanced, but neonatal morbidity was not increased. This may be related to early intervention in the form of cesarean or instrumental delivery. Fetal distress, non-reassuring fetal status, fetal growth restriction and post maturity are the factors which can lead to meconium

aspiration syndrome¹³. A prospective observational study carried out in Shillong, India by Mundhra and Aggarwal² pointed out the worrisome complications of meconium in amniotic fluid in gestations beyond term. They included 165 cases of term pregnancies with meconium stained liquor and compared with 190 controls. Fifty percent cases were of more than 40 weeks of gestation. Cesarean rate was found to be double to that of control population with clear liquor (81% vs. 49%). There was increased rate of neonatal admissions, birth asphyxia and meconium aspiration syndrome. Post term pregnancies have been associated with varied complications like increased frequency of neonatal convulsions, meconium aspiration syndrome and Apgar score of less than 4 at 5 minutes¹⁵. In our study also, cesarean rate was found to be very high in pregnancies with MSL, however neonatal morbidity was not seen.

In our study, 42.8% women were induced mainly for postdated pregnancy. In a study by Kassis et al¹⁵ done in Israel, postdated pregnancies had higher rate of induction (40% vs. 3%) and meconium stained liquor (31 vs 15.5%). The relative risk of fetal distress in the form of non-reassuring cardiotocography in women with meconium stained liquor increased with more advanced gestational age¹⁶. Hence, there is a higher rate of induction in postdated pregnancies.

In our institute, incidence of cesarean section is high in pregnancies with MSL. The most common reason is a poor bishop's score at admission, which leads to a decision of not giving a trial of induction. In our institute in the absence of provision of fetal scalp blood monitoring, non-reassuring features in cardiotocograph are considered as an indication of cesarean section in these pregnancies, especially if discovered in early labor.

This was a retrospective study with a relatively small sample of postdates women. Prevalence of MSL in term pregnancies before 40 weeks and its comparison with those beyond 40 weeks is not included in this study. A prospective multicentric study with longer follow up of the neonate would further elucidate the chances of meconium passage at different gestations beyond term and the odds of neonatal morbidity and mortality. Since prevalence of MSL is high in our fetuses after 40 weeks and increases with increasing gestational age, prospective studies should be done to make Indian guidelines for early induction of labour after 40 weeks.

Conclusion

The prevalence of meconium stained liquor in Indian women at more than 40 weeks was 14.92% at our institute. Prevalence increases with increasing gestational age. Most of the pregnancies with MSL eventually delivered via cesarean section. Neonatal outcome is favourable in babies born with MSL.

Conflict of interest: None. **Disclaimer:** Nil.

References

1. Kalra VK, Lee HC, Sie L, Ratnasiri AW, Underwood MA, Lakshminrusimha S. Change in neonatal resuscitation guidelines and trends in incidence of meconium aspiration syndrome in California. *J Perinatol.* 2020 Jan; 40(1): 46-55.
2. Mundhra R, Aggarwal M. Fetal outcome in meconium stained deliveries. *J Clin Diagn Res JCDR.* 2013 Dec; 7(12): 2874-6.
3. Caughey AB, Musci TJ. Complications of term pregnancies beyond 37 weeks of gestation. *Obstet Gynecol.* 2004 Jan;103(1): 57-62.
4. Caughey AB, Sundaram V, Kaimal AJ, Cheng YW, Gienger A, Little SE, et al. Maternal and neonatal outcomes of elective induction of labor. *Evid Report Technology Assess.* 2009 Mar; 176: 1-257.
5. American College of Obstetricians and Gynecologists. Practice bulletin no. 146: Management of late-term and postterm pregnancies. *Obstet Gynecol.* 2014;124 (2 Pt 1): 390-6.
6. Hamilton BE, Martin JA, Osterman MJ, Curtin SC, Matthews TJ. Births: Final Data for 2014. *Natl Vital Stat Rep.* 2015 Dec; 64(12):1-64.
7. Caughey AB, Stotland NE, Washington AE, Escobar GJ. Who is at risk for prolonged and postterm pregnancy? *Am J Obstet Gynecol.* 2009 Jun; 200(6): 683.e1-5.
8. Weiss E, Abele H, Bartz C, Franz M, Fischer T, Gembruch U, et al. S1-Guideline: Management of Late-term and Post-term Pregnancy. *Geburtshilfe Frauenheilkd.* 2014 Dec; 74(12):1099-103.
9. Sedaghatian MR, Othman L, Hossain MM, Vidyasagar D. Risk of meconium-stained amniotic fluid in different ethnic groups. *J Perinatol Off J Calif Perinat Assoc.* 2000 Jun; 20(4): 257-61.
10. Narasimhaiah A, Pratibha SD. Mode of delivery and foetal outcome in meconium-stained liquor: a

- retrospective study. *Journal of Evidence Based Medicine and Healthcar.* 2016 Aug 2;3(62):3342-4.
11. Addisu D, Asres A, Gedefaw G, Asmer S. Prevalence of meconium stained amniotic fluid and its associated factors among women who gave birth at term in Felege Hiwot comprehensive specialized referral hospital, North West Ethiopia: a facility based cross-sectional study. *BMC Pregnancy Childbirth.* 2018 Oct 30; 18(1): 429.
 12. Hiersch L, Krispin E, Linder N, Aviram A, Gabbay-Benziv R, Yogev Y, et al. Meconium-Stained Amniotic Fluid and Neonatal Morbidity in Low-Risk Pregnancies at Term: The Effect of Gestational Age. *Am J Perinatol.* 2017; 34(2):183-90.
 13. Chand S, Salman A, Abbassi RM, Siyal AR, Ahmed F, Leghari AL, et al. Factors Leading To Meconium Aspiration Syndrome in Term- and Post-term Neonates. *Cureus.* 2019 Sep 5;11(9):e5574.
 14. Galal M, Symonds I, Murray H, Petraglia F, Smith R. Postterm pregnancy. *Facts Views Vis ObGyn.* 2012; 4(3):175-87.
 15. Kassis A, Mazor M, Leiberman JR, Cohen A, Insler V. Management of post-date pregnancy: a case control study. *Isr J Med Sci.* 1991 Feb; 27(2): 82-6.
 16. Wong SF, Chow KM, Ho LC. The relative risk of “fetal distress” in pregnancy associated with meconium-stained liquor at different gestation. *J Obstet Gynaecol.* 2002 Jan 1; 22(6): 594-9.

Shatakshi Kulshreshtha¹, Sujata Siwatch²

¹ Senior Resident, Department of Obstetrics and Gynecology, PGIMER Chandigarh, India; ² Associate Professor, Department of Obstetrics and Gynecology, PGIMER Chandigarh, India.